

# On the construction of set-valued dual processes

Laurent Miclo

Institut de Mathématiques de Toulouse  
École d'Économie de Toulouse

# Plan of the talk

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion
- 6 Some references

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion
- 6 Some references

# Convergence to equilibrium

Consider  $X := (X_t)_{t \geq 0}$  an ergodic Markov process on some state space  $V$  and let  $\pi$  be its invariant probability measure.

In practice,  $\pi$  is given and  $X$  is typically constructed via a Metropolis-type procedure.

Our obsession: the quantitative convergence to equilibrium.

The deterministic **mixing time**

$$T_\epsilon := \inf\{t \geq 0 : \|\mathcal{L}(X_t) - \pi\|_{\text{tv}} \leq \epsilon\}$$

enables to stop the algorithm  $X$  to get a r.v.  $X_{T_\epsilon}$  sampled according to  $\pi$ , up to the precision  $\epsilon \in (0, 1)$ .

# Strong stationary times (1)

Strong stationary times: a probabilistic approach to convergence to equilibrium: by looking at a given trajectory, one decides to stop it at a random time to get an *exact* sample of  $\pi$ .

A **strong stationary time**  $\tau$  associated to  $X$  is a finite stopping time such that

$$\tau \perp\!\!\!\perp X_\tau \quad \text{and} \quad X_\tau \sim \pi$$

# Strong stationary times (2)

Leads to estimates on the speed of convergence:

$$\forall t \geq 0, \quad \|\mathcal{L}(X_t) - \pi\|_{\text{tv}} \leq \mathfrak{s}(\mathcal{L}(X_t), \mu) \leq \mathbb{P}[\tau > t]$$

in **total variation** and in **separation discrepancy**: for any probability measures  $\mu$  and  $\pi$  on the same state space:

$$\mathfrak{s}(\mu, \pi) := \text{esssup}_{\pi} 1 - \frac{d\mu}{d\pi} \geq \frac{1}{2} \left\| \frac{d\mu}{d\pi} - 1 \right\|_{\mathbb{L}^1(\pi)} =: \|\mu - \pi\|_{\text{tv}}$$

(when  $\mu \ll \pi$  for the last equality).

Strong stationary times were introduced by [Aldous and Diaconis \[1986\]](#) to investigate the quantitative convergence to equilibrium of the top-to-random card shuffle.

# Markov intertwining relations (1)

How to obtain a strong stationary time?

Find an absorbed **dual** Markov process  $\mathfrak{X} := (\mathfrak{X}_t)_{t \geq 0}$  on a state space  $\mathfrak{V}$  such that there exists  $\Lambda$  a Markov kernel from  $\mathfrak{V}$  to  $V$  satisfying the **intertwining relations**

$$\begin{aligned}\mathcal{L}(\mathfrak{X}_0)\Lambda &= \mathcal{L}(X_0) \\ \mathfrak{L}\Lambda &= \Lambda L\end{aligned}$$

where  $L$  and  $\mathfrak{L}$  are the generators of  $X$  and  $\mathfrak{X}$ .

## Markov intertwining relations (2)

Then there is a coupling of  $X$  and  $\mathfrak{X}$  such that the absorption time for  $\mathfrak{X}$  is a strong stationary time for  $X$ .

This method was developed by [Diaconis and Fill \[1990\]](#), at least for discrete time and finite state spaces  $V$  and  $\mathfrak{V}$ . The coupling was such that, for all time  $n \in \mathbb{Z}_+$ ,

$$\begin{aligned}\mathcal{L}(\mathfrak{X}_{[0,n]}|X) &= \mathcal{L}(\mathfrak{X}_{[0,n]}|X_{[0,n]}) \\ \mathcal{L}(X_n|\mathfrak{X}_{[0,n]}) &= \Lambda(\mathfrak{X}_n, \cdot)\end{aligned}$$

# Set-valued dual processes

An interesting class of absorbed dual processes are those which are **set-valued**:  $\mathfrak{V}$  is a nice set of measurable subsets  $S$  of  $\mathfrak{X}$  such that  $\pi(S) > 0$  or  $S$  is a singleton. The kernel  $\Lambda$  corresponds to the **conditional expectation** under  $\pi$ : for any  $S \in \mathfrak{V}$ ,

$$\Lambda(S, \cdot) = \begin{cases} \frac{\pi(S \cap \cdot)}{\pi(S)} & , \text{ if } \pi(S) > 0 \\ \delta_x & , \text{ if } S = \{x\} \end{cases}$$

Furthermore, the process  $\mathfrak{X}$  is assumed to be absorbed at  $V \in \mathfrak{V}$ .

# Pitman's theorem

A famous example is **Pitman's intertwining relation** between the Brownian motion and the Bessel process [Pitman 1975, Pitman and Rogers 1981]. Here  $V = \mathbb{R}$ ,  $X$  is the Brownian motion starting from 0 and

$$\mathfrak{V} := \{[-r, r] : r \geq 0\}$$

The dual process  $\mathfrak{X} := ([-R_t, R_t])_{t \geq 0}$  is given by

$$\forall t \geq 0, \quad R_t := 2M_t^X - X_t$$

where  $M^X := (M_t^X)_{t \geq 0}$  is the **maximum process**:

$$\forall t \geq 0, \quad M_t^X := \max\{X_s : s \in [0, t]\}$$

The process  $(R_t)_{t \geq 0}$  is known to be a **Bessel-3 process**, namely has the same law as the norm of a Brownian motion in dimension 3.

# Pitman's theorem in picture




Figure: Trajectories: Brownian motion  $B_{[0,t]}$ ,  $R_{[0,t]}$ ,  $-R_{[0,t]}$ , and the segment-valued dual:  $[-R_t, R_t]$

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion
- 6 Some references

# Random mappings (1)

Consider the discrete time and finite state space situation. Let  $P := (P(x, y))_{x, y \in V}$  be the transition matrix of  $X := (X_n)_{n \in \mathbb{Z}_+}$ . Assume  $P$  irreducible and let  $\pi$  be the invariant probability measure. The **adjoint transition matrix**  $P^* := (P^*(x, y))$  is given by

$$\forall x, y \in V, \quad P^*(x, y) := \frac{\pi(y)}{\pi(x)} P(y, x)$$

A random mapping  $\psi : V \rightarrow V$  is said to be **associated to  $P^*$**  when

$$\forall x, x' \in V, \quad \mathbb{P}[\psi(x) = x'] = P^*(x, x')$$

It is convenient to have at our disposal a family of random mappings  $(\psi_S)_{S \in \tilde{\mathfrak{M}}}$  associated to  $P^*$ , where  $\tilde{\mathfrak{M}} := \{A : A \subset V\}$ .

## Random mappings (2)

Such a family enables to define a random mapping  $\Psi$  from  $\bar{\mathfrak{Y}}$  to  $\bar{\mathfrak{Y}}$  via

$$\forall S \in \bar{\mathfrak{Y}}, \quad \Psi(S) := \{y \in V : \psi_S(y) \in S\}$$

Consider the transition matrix  $K$  from  $\bar{\mathfrak{Y}}$  to  $\bar{\mathfrak{Y}}$  given by

$$\forall S, S' \in \bar{\mathfrak{Y}}, \quad K(S, S') := \mathbb{P}[\Psi(S) = S']$$

as well as its **Doob transform**:

$$\forall S, S' \in \mathfrak{Y}, \quad \mathfrak{P}(S, S') := \frac{\pi(S')}{\pi(S)} K(S, S')$$

where  $\mathfrak{Y} = \bar{\mathfrak{Y}} \setminus \{\emptyset\}$ .

# Conditioned random mappings (1)

For  $x, x' \in V$  with  $P(x, x') > 0$  and  $S \in \mathfrak{V}$  containing  $x$ , denote

$$\forall S' \in \mathfrak{V}, \quad K_{x, x'}(S, S') := \mathbb{P}[\Psi(S) = S' | \psi_S(x') = x]$$

Note that the conditioning is non-degenerate, since

$$\mathbb{P}[\psi_S(x') = x] = P^*(x', x) > 0$$

Consider

$$W := \{(x, S) \in V \times \mathfrak{V} : x \in S\}$$

## Conditioned random mappings (2)

Let  $\mathcal{A}$  be the set of probability measures  $m$  on  $W$  of the form

$$\forall (x, S) \in W, \quad m(x, S) = \mu(S)\Lambda(S, x)$$

where  $\mu$  is the marginal of  $m$  on  $\mathfrak{X}$ .

Define a Markov kernel  $Q$  on  $W$  via

$$\begin{aligned} \forall (x, S), (x', S') \in W \\ Q((x, S), (x', S')) := P(x, x')K_{x, x'}(S, S') \end{aligned}$$

## Theorem 1

*Let  $(X_n, \mathfrak{X}_n)_{n \in \mathbb{Z}_+}$  be a Markov chain on  $W$  whose initial distribution  $\mathcal{L}(X_0, \mathfrak{X}_0)$  belongs to  $\mathcal{A}$  and whose transitions are given by  $Q$ . Then  $\mathfrak{X} := (\mathfrak{X}_n)_{n \in \mathbb{N}}$  is a set-valued absorbed dual for  $X := (X_n)_{n \in \mathbb{N}}$ . Furthermore,  $\mathfrak{X}$  is Markovian and its transitions are given by  $\mathfrak{P}$ .*

This result is related to the coupling-from-the-past algorithm of [Propp and Wilson \[1996\]](#) and to the evolving set process of [Morris and Peres \[2005\]](#). There is an improvement based on random mappings weakly associated to  $P^*$ .

# A corresponding algorithm

Let be given a trajectory  $(x_n)_{n \in \mathbb{Z}_+}$  of  $X$ .

- Start with  $\mathfrak{X}_0 := \{x_0\}$ .
- When  $\mathfrak{X}_n$  has been constructed, consider a random mapping  $\psi_{\mathfrak{X}_n}$  (weakly) associated to  $P^*$ , whose law may depend on  $\mathfrak{X}_n$ .
- Condition by the fact that  $\psi_{\mathfrak{X}_n}(x_{n+1}) = x_n$  and sample a corresponding mapping  $\varphi$ , to construct

$$\mathfrak{X}_{n+1} := \{y \in V : \varphi(y) \in \mathfrak{X}_n\}$$

(due to the conditioning, we have  $x_{n+1} \in \mathfrak{X}_{n+1}$ ).

This works for any random mapping associated to  $P^*$ , all the difficulty stays on relevant choices leading to (close to optimal) strong stationary times.

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem**
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion
- 6 Some references

# A denumerable state space example (1)

Consider the transition “matrix” of the simple random walk on  $\mathbb{Z}$ :

$$\forall x, y \in \mathbb{Z}, \quad P(x, y) := \begin{cases} 1/2 & , \text{ if } |y - x| = 1 \\ 0 & , \text{ otherwise} \end{cases}$$

An invariant measure  $\pi$  is the counting measure, it is even reversible for  $P$ , in the sense that  $P^* = P$ .

The kernel  $\Lambda$  is still well-defined, if we take for  $\mathfrak{V}$  the set of finite non-empty subsets of  $\mathbb{Z}$ .

## A denumerable state space example (2)

Let  $X := (X_n)_{n \in \mathbb{Z}_+}$  be a random walk with transition kernel  $P$  and starting from 0.

Introduce the process  $R^\vee := (R_n^\vee)_{n \in \mathbb{Z}_+}$  defined by

$$\forall n \in \mathbb{Z}_+, \quad R_n^\vee := 2 \max\{X_m : m \in \llbracket 0, n \rrbracket\} - X_n$$

Finally consider  $\mathfrak{X} := (\mathfrak{X}_n)_{n \in \mathbb{Z}_+}$  given by

$$\forall n \in \mathbb{Z}_+, \quad \mathfrak{X}_n := \{R_n^\vee - 2m : m \in \llbracket 0, R_n^\vee \rrbracket\}$$

[Pitman \[1975\]](#) has shown that  $\mathfrak{X}$  is a set-valued dual for  $X$ .

# A corresponding random mapping

Consider the function  $\psi$  given by

$$\forall S \in \mathfrak{V}, \forall x \in \mathbb{Z}, \forall b \in \{-1, 1\},$$
$$\psi(S, x, b) := \begin{cases} x + b & , \text{ if } x > \max(S) \\ x - b & , \text{ if } x \leq \max(S) \end{cases}$$

Consider a Rademacher variable  $B$ , i.e. such that

$\mathbb{P}[B = -1] = \mathbb{P}[B = 1] = 1/2$  and for fixed  $S \in \mathfrak{V}$ , let  $\psi_S$  be the random mapping given by

$$\forall x \in \mathbb{Z}, \quad \psi_S(x) := \psi(S, x, B)$$

It is clear that  $\psi_S$  is a random mapping associated to  $P^* = P$ . The discrete Pitman's theorem can be easily deduced from Theorem 1 with the family  $(\psi_S)_{S \in \mathfrak{V}}$ .

# Schematic proof




Figure: Schematic proof of the discrete Pitman theorem via random mappings

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle**
- 5 Exemple 3: Brownian motion
- 6 Some references

# A card shuffle example (1)

The **top-to-random shuffle** takes the top card of a deck and put it at a uniform random location inside. With  $N$  the number of cards, it leads to a Markov chain  $X := (X_n)_{n \in \mathbb{Z}_+}$  on the symmetric group  $V := \mathcal{S}_N$ , starting from id and whose transition matrix  $P := (P(\sigma, \sigma'))_{\sigma, \sigma' \in \mathcal{S}_N}$  is given by

$$P(\sigma, \sigma') := \begin{cases} 1/N & , \text{ if there exists } l \in \llbracket 1, N \rrbracket \text{ with} \\ & \sigma' = (1 \rightarrow l \rightarrow l-1 \rightarrow \dots \rightarrow 2) \circ \sigma \\ 0 & , \text{ otherwise} \end{cases}$$

$P$  is irreducible and its invariant probability  $\pi$  is the uniform probability distribution on  $\mathcal{S}_N$ .

## A card shuffle example (2)

The Markov chain  $X$  admits a famous set-valued dual process  $\tilde{\mathfrak{X}} := (\tilde{\mathfrak{X}}_n)_{n \in \mathbb{Z}_+}$  defined by [Aldous and Diaconis \[1986\]](#):

$$\forall n \in \mathbb{Z}_+, \quad \tilde{\mathfrak{X}}_n := A_{X_n, Y_n}$$

where  $Y_n \in \llbracket 1, N \rrbracket$  is the position of the initial last card and for any  $\sigma \in \mathcal{S}_N$  and  $y \in \llbracket 0, N \rrbracket$ ,

$$A_{\sigma, y} := \{\sigma' \in \mathcal{S}_N : \sigma'(1) = \sigma(1), \dots, \sigma'(y) = \sigma(y)\}$$

The associated strong stationary time  $\tilde{\tau}$  is first time the initial last card arrives at the top of the deck and is inserted. It is easy to check that  $\mathbb{E}[\tilde{\tau}] \sim N \ln(N)$ .

# A corresponding random mapping (1)

Consider the first time  $\tau$  the initial last-but-one card arrives at the top of the deck and is inserted.

Let us check via random mappings that  $\tau$  is also a strong stationary time, strictly better than  $\tilde{\tau}$ .

Here  $P^*$  is the transition matrix of the **random-to-top shuffle**.

Consider for any  $x \in \llbracket 1, N \rrbracket$ , the mapping  $\psi^{(x)} : \mathcal{S}_N \rightarrow \mathcal{S}_N$  which acts on any permutation  $\sigma$  by removing the card  $x$  from the deck and putting it at the top. Formally, we have

$$\forall \sigma \in \mathcal{S}_N, \quad \psi^{(x)}(\sigma) = (1 \rightarrow 2 \rightarrow \dots \rightarrow \sigma^{-1}(x)) \circ \sigma$$

( $\sigma^{-1}(x)$  is the position of the card  $x$ ).

## A corresponding random mapping (2)

Let  $(U_n)_{n \in \mathbb{N}}$  be a family of independent random variables uniformly distributed on  $\llbracket 1, N \rrbracket$ . At any time  $n \in \mathbb{N}$ , consider the random mapping  $\psi^{(U_n)}$ , it is associated to  $P^*$ . Here there is no dependence on a subset  $S \in \mathfrak{V}$ .

Let be given a trajectory  $x_{\llbracket 0, n \rrbracket}$ , for some fixed  $n \in \mathbb{Z}_+$ , starting from the identity,  $x_0 = \text{id}$ . For any  $m \in \llbracket 1, n \rrbracket$ , let  $\varphi_m$  be the conditioning of  $\psi^{(U_m)}$  by  $\psi^{(U_m)}(x_m) = x_{m-1}$ . As in the previous example,  $\varphi_m$  is deterministic, as we have  $\varphi_m = \psi^{(x_{m-1}(1))}$ .

We have

$$\forall n \in \mathbb{Z}_+, \quad \mathfrak{X}_n = \{\sigma \in \mathcal{S}_N : \varphi_1 \circ \varphi_2 \circ \cdots \circ \varphi_n(\sigma) = \text{id}\}$$

and the corresponding absorption time at  $\mathcal{S}_N$  is  $\tau$ .

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion**
- 6 Some references

Theorem 1 admits several extensions:

- the state space  $V$  can be a Polish space,
- the time can be continuous (for the moment, only in a regular diffusion framework).

The underlying theory is quite involved and lacks the existence of convenient stochastic coalescing flows depending on subset evolutions, despite the works of [Le Jan and Raimond \[2004, 2006\]](#).

An illustration of the use of stochastic coalescing flows on the classical Brownian motion:

### Theorem 2

*The process  $\mathfrak{X} = ([-(L_t^0(X) + |X_t|), L_t^0(X) + |X_t|)]_{t \geq 0}$  is a set-valued dual for the Brownian motion  $X$  starting from 0, where  $L^0(X) := (L_t^0(X))_{t \geq 0}$  is the local time of  $X$  at 0.*

One recover Pitman's theorem, by taking into account Lévy's theorem stating that for any  $t \geq 0$

$$(M_t - X_t, M_t)_{t \geq 0} \stackrel{\mathcal{L}}{=} (|X_t|, L_t^0(X))_{t \geq 0}$$

# Stochastic coalescing flows (1)

Let  $B := (B_s)_{s \geq 0}$  be another Brownian motion. Consider the following system of Tanaka's equations, for all  $t \geq 0$  and  $y \in \mathbb{R}$ ,

$$\begin{cases} dY_s^{(t)}(y) &= -\text{sgn}(Y_s^{(t)}(y))dB_s^{(t)}, & \forall s \in [0, t] \\ Y_0^{(t)}(y) &= y \end{cases} \quad (1)$$

where  $\text{sgn}$  equals  $-1$  on  $(-\infty, 0]$  and  $1$  on  $(0, +\infty)$  and where  $B^{(t)} := (B_{t-s})_{s \in [0, t]}$  is the time-reversed process associated to  $B$  at time  $t \geq 0$ .

# Stochastic coalescing flows (2)

Le Jan and Raimond [2006] provide a (non-Wiener) coalescing stochastic flow solution to this system.

Define  $\psi := (\psi_{s,t}(y))_{(s,t,y) \in \Delta \times \mathbb{R}}$  via

$$\forall x \in \mathbb{R}, \forall 0 \leq s \leq t, \quad \psi_{s,t}(y) := Y_{t-s}^{(t)}(y)$$

with  $\Delta := \{(s, t) : 0 \leq s \leq t\}$ .

By monotonicity in  $y$ , there is a version of  $\psi$  which is càdlàg in  $y$ .

# Conditioned stochastic coalescing flows (1)

Fix  $t \geq 0$  and a Brownian trajectory  $X_{[0,t]}$ . Conditioning  $\psi$  by the event

$$\forall s \in [0, t], \quad \psi_{s,t}(X_t) = X_s$$

implies in particular that  $X^{(t)}$  is a solution to Tanaka's stochastic differential equation:

$$\forall s \in [0, t], \quad dX_s^{(t)} = -\operatorname{sgn}(X_s^{(t)})dB_s^{(t)}$$

namely the white noise associated to  $B^{(t)}$  is determined by  $X^{(t)}$ :

$$\forall s \in [0, t], \quad dB_s^{(t)} = -\operatorname{sgn}(X_s^{(t)})dX_s^{(t)}$$

# Conditioned stochastic coalescing flows (2)

We deduce that the conditioned flow  $\varphi$  is given by

$$\forall 0 \leq s \leq t, \forall z \in \mathbb{R}, \quad \varphi_{s,t}(z) := Z_s^{(t)}(z)$$

where

$$\begin{cases} dZ_s^{(t)}(z) &= \operatorname{sgn}(Z_s^{(t)}(z)) \operatorname{sgn}(X_s^{(t)}) dX_s^{(t)} \\ Z_0^{(t)}(z) &= z \end{cases}$$

This system is the same as (1), once  $B^{(t)}$  is replaced by the Brownian motion  $(-\int_0^s \operatorname{sgn}(X_v^{(t)}) dX_v^{(t)})_{s \in [0,t]}$ .

# Conditioned stochastic coalescing flows (3)

Theorem 2 is deduced from the fact that  $\mathfrak{X} := (\mathfrak{X}_t)_{t \geq 0}$ , defined by

$$\forall t \geq 0, \quad \mathfrak{X}_t := \varphi_{0,t}^{-1}(\{0\})$$

is a set-valued dual for  $X$  and that we can compute

$$\forall t > 0, \quad \mathfrak{X}_t = [-(L_t^0(X) + |X_t|), L_t^0(X) + |X_t|]$$

In the underlying theory, one has to consider

$$\mathfrak{V} := \{[a, b) : a < b \in \mathbb{R}\}$$

to which is added the initial subset  $\mathfrak{X}(0) = \{0\}$ .

# What we really want (1)

We would like to have at our disposal a solution to the following system of equations, for all  $0 \leq s \leq t$  and  $y \in \mathbb{R}$ ,

$$\begin{cases} dY_s^{(t)}(y) &= \operatorname{sgn}(R_{t-s}^\vee - Y_s^{(t)}(y))dB_s^{(t)} + b(Y_s^{(t)}(y))ds \\ Y_0^{(t)}(y) &= y \\ R_{t-s}^\vee &:= \max\{z \in \mathbb{R} : Y_{t-s}^{(t-s)}(z) \in \mathfrak{X}_0\} \end{cases}$$

where  $b : \mathbb{R} \rightarrow \mathbb{R}$  is a nice drift.

## What we really want (2)

For  $b = 0$ , it would lead to a proof of Pitman's theorem similar to that given in the discrete situation. Above all, for  $b \neq 0$ , it would provide a direct coupling of the diffusion  $X$  solution of the s.d.e.

$$\forall t \geq 0, \quad dX_t = dW_t + b(X_t)dt$$

(where  $W := (W_t)_{t \geq 0}$  is a Brownian motion) with a non-trivial segment-valued dual process. It would open the way for multidimensional and hypo-elliptic extensions, which are the remote motivation for this work.

- 1 Introduction
- 2 Construction via random mappings
- 3 Exemple 1: discrete Pitman's theorem
- 4 Exemple 2: top-to-random shuffle
- 5 Exemple 3: Brownian motion
- 6 Some references**

# Primary references on intertwining relations



James W. Pitman.

One-dimensional Brownian motion and the three-dimensional Bessel process.

*Advances in Appl. Probability*, 7(3):511–526, 1975.



L. Chris G. Rogers and James W. Pitman.

Markov functions.

*Ann. Probab.*, 9(4):573–582, 1981.



David Aldous and Persi Diaconis.

Shuffling cards and stopping times.

*Amer. Math. Monthly*, 93(5):333–348, 1986.



Persi Diaconis and James Allen Fill.

Strong stationary times via a new form of duality.

*Ann. Probab.*, 18(4):1483–1522, 1990.



David A. Levin, Yuval Peres, and Elizabeth L. Wilmer.

*Markov chains and mixing times*.

American Mathematical Society, Providence, RI, 2009.

# References on some *geometric* processes



James Gary Propp and David Bruce Wilson.

Exact sampling with coupled Markov chains and applications to statistical mechanics.

In *Proceedings of the Seventh International Conference on Random Structures and Algorithms (Atlanta, GA, 1995)*, volume 9, pages 223–252, 1996.



Ben Morris and Yuval Peres.

Evolving sets, mixing and heat kernel bounds.

*Probab. Theory Related Fields*, 133(2):245–266, 2005.



Yves Le Jan and Olivier Raimond.

Flows, coalescence and noise.

*Ann. Probab.*, 32(2):1247–1315, 2004.



Yves Le Jan and Olivier Raimond.

Flows associated to Tanaka's SDE.

*ALEA Lat. Am. J. Probab. Math. Stat.*, 1:21–34, 2006.

# Further references on Markov intertwining



Philippe Carmona, Frédérique Petit and Marc Yor.

Beta-gamma random variables and intertwining relations between certain Markov processes.

*Rev. Mat. Iberoamericana*, 14(2):311–367, 1998.



Alexei Borodin and Grigori Olshanski.

Markov processes on the path space of the Gelfand-Tsetlin graph and on its boundary.

*J. Funct. Anal.*, 263(1):248–303, 2012.



Soumik Pal and Mykhaylo Shkolnikov.

Intertwining diffusions and wave equations.

*ArXiv e-prints*, June 2013.



Laurent Miclo.

On the construction of set-valued dual processes.

*HAL e-prints*, November 2018.